

Clinical Consideration of Obese Infertile Women

Yoon Sung Nam, Chang Jin Jeong, Nam Keun Kim^{*}, Tae Ki Yoon, Kwang Yul Cha

Department of Obstetrics and Gynecology, Institute for Clinical Research^{},
College of Medicine, Pocheon CHA University, Pocheon, Korea*

Objective: To analyze the clinical characteristics of obese infertile women.

Material and Method: Height, weight, body mass index, menstrual pattern, glucose, insulin, glucose / insulin ratio, dehydroepiandrosterone sulfate (DHEA -S), testosterone, free testosterone and plasminogen activator inhibitor (PAI-1) of 15 obese infertile women were tested.

Results: Of 15 obese infertile women, the number of diabetes mellitus, hyperinsulinemia, and insulin resistance was 2 (13%), 2 (13%), 2 (13%), respectively. The incidence of increased DHEA -S, testosterone, and free testosterone was 7 (47%), 1 (7%), 6 (40%), respectively. Notably, all patients showed increased PAI-1.

Conclusions: Obesity is associated with infertility as well as many kinds of health problems. Obesity is closely related to insulin resistance and it also causes hyperandrogenism. Increased PAI-1 is one of the important causes of thrombophilia. Consequently, in the workup of obese infertile patient, many aspects of health problems should be considered.

Key Words: Obesity, Infertility

(obesity) (visceral obesity) (insulin resistance)
(adiposity) (hyperinsulinemia)
(overweightness)
(sex hormone binding globulin: SHBG)
testosterone 가
가 가 (body mass index: (clearance rate) 가
BMI) (kg) (m)
가 25 , (aromatization) 가
30
(upper body obesity)

: ,) 135-081 1 650-9,
Tel: (02) 3468-3412, Fax: (02) 555-9007, e-mail: ysnam@cha.ac.kr

Table 1. Clinical characteristics of obese infertile women

	Height	Weight	BMI	Glucose	Insulin	G / I	DS	T	Free T	PAI-1
1	159	71	28.1	91	3.1	29.3	N	N		
2	152	70	30.4	88	9.4	9.4	N	N	N	
3	160	76	29.7	81	1.0	81.0	N		N	
4	157	74	30.1	294	5.6	52.5		N	N	
5	162	73	27.9	83	26.7	3.1				
6	158	70	28.0	91	10.8	8.43	N	N	N	
7	161	90	34.7	136	32.8	4.15		N		
8	167	92	33.0	85	16.0	5.3		N		
9	171	75	25.7	118	7.7	15.3		N	N	
10	168	73	25.9	93	16.1	5.8	N	N		
11	158	78	31.2	100	6.5	15.4	N	N	N	
12	157	82	33.3	111	15.1	7.4	N	N		
13	163	70	26.3	81	1.0	81.0		N	N	
14	160	82	32.0	83	5.3	15.7	N	N	N	
15	158	85	34.0	93	4.7	19.8		N	N	

BMI: body mass index, G / I: glucose / insulin, DS: dehydroepiandrosterone sulfate, T: testosterone, PAI-1: plasminogen activator inhibitor-1

factor 1: IGF-1)
 가
 1 (insulin-like growth factor)
 leptin
 가
 1.
 2001 1 1 2001 12 31
 가
 25 15
 가 가
 2.
 가
 (hyperlipidemia) 가
 (glucose intolerance) 가
 , dehydroepiandrosterone sulfate (DHEA-S), testosterone, testosterone, plasminogen activator inhibitor (PAI-1)
 가
 가
 15 가 2 (13%)

sterone 1 (7%),
 15 가

2 (13%)
 2 (13%)
 DHEA-S 7 (47%), testosterone 6 (40%)
 PAI-1 (Table 1).

가 (gonadotropin releasing hormone: GnRH) (agonist) 6

가 (hyperandrogenism), (hyperinsulinemia) 10,11

가 (upper body obesity) 1 가 (insulin resistance)

가 (gynecoid obesity) (waist-to-hip ratio) 가 (sex hormone binding globulin: SHBG) 가 가 가 2 가 3 가 13,14 가 4 가 5 가 6 가 16,17 가 7 (euglycemic hyperinsulinemic clamp procedure) testosterone 8 가 19 (antiandrogen)

PAI-1 가

가

가

가

1. Dunaif A, Futterweit W, Segal KR, Dobrjansky A. Profound peripheral insulin resistance, independent of obesity, in the polycystic ovary syndrome. *Diabetes* 1989; 38: 1165-74.
2. Meirow D, Yossepowitch O, Rosler A. Insulin resistant and non-resistant polycystic ovary syndrome present two clinical and endocrinologic subgroups. *Hum Reprod* 1995; 10: 1951-6.
3. Evans DJ, Hoffman RG, Kalkoff RK, Kissebah AH. Relationship of androgen activity to body fat topography, fat cell morphology, and metabolic aberrations in premenopausal women. *J Clin Endocrinol Metab* 1983; 57: 304-10.
4. Ohlson LO, Larsson B, Svardsudd K. The influence of body fat distribution on the incidence of diabetes mellitus. *Diabetes* 1985; 34: 1055-8.
5. Flier JS, Eastman RC, Minaker KL, Matteson D, Rowe JW. Acanthosis nigricans in obese women with hyperandrogenism: Characterization of an insulin-resistant state distinct from the type A and B syndromes. *Diabetes* 1985; 34: 101-7.
6. Kissebah AH, Vydellingum N, Murray R. Relation of body fat distribution to metabolic complications of obesity. *J Clin Endocrinol Metab* 1982; 54: 254-60.
7. Billar RB, Rishardson D, Schwartz R, Posner B, Little B. Effect of chronically elevated androgen or estrogen on the glucose tolerance test and insulin response in female rhesus monkeys. *Am J Obstet Gynecol* 1987; 157: 1297-302.
8. Holmang A, Larsson BM, Brzezinska Z, Bjorntorp P. Effects of short-term testosterone exposure on insulin sensitivity of muscles in female rats. *Am J Physiol* 1992; 25: 851-5.
9. Moghetti P, Tosi F, Castello R. The insulin resistance in women with hyperandrogenism is partially reversed by antiandrogen treatment: evidence that androgens impair insulin action in women. *J Clin Endocrinol Metab* 1996; 81: 952-60.
10. Dunaif A, Green G, Futterweit W. Suppression of hyperandrogenism does not improve peripheral or hepatic insulin resistance in the PCOS. *J Clin Endocrinol Metab* 1990; 70: 699-704.
11. Geffner ME, Kaplan SA, Bersch N, Golde DW, Landaw EM, Chang RJ. Persistence of insulin resistance in polycystic ovarian disease after inhibition of ovarian steroid secretion. *Fertil Steril* 1986; 45: 327-33.
12. Dunaif A. Insulin resistance and the polycystic ovary syndrome: mechanism and implications for pathogenesis. *Endocr Rev* 1997; 18: 774-800.
13. Stuart CA, Prince MJ, Peters EJ, Meyer WJ. Hyperinsulinemia and hyperandrogenemia: in vivo androgen response to insulin infusion. *Obstet Gynecol* 1987; 69: 921-5.
14. Stuart CA, Nagamani N. Insulin infusion acutely augments ovarian androgen production in normal women. *Fertil Steril* 1990; 54: 788-92.
15. Nestler JE, Barlascini CO, Matt DW. Suppression of serum insulin by diazoxide reduces serum testosterone levels in obese women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 1989; 68: 1027-32.
16. Nestler JE, Strauss JF. Insulin as an effector of human ovarian and adrenal steroid metabolism. *Endocr Metab Clin North Am* 1991; 20: 807-23.
17. Dunaif A. Insulin resistance and ovarian hyperandrogenism. *Endocrinologist* 1992; 2: 248-60.
18. Barbieri RL, Makris A, Randall RW. Insulin stimulates androgen accumulation in incubations of ovarian stroma obtained from women with hyperandrogenism. *J Clin Endocrinol Metab* 1986; 62:

- 904-10.
19. Plymate SR, Matej LA, Jones RE, Freidl KE. Inhibition of sex hormone-binding globulin production in the human hepatoma (Hep G2) cell line by insulin and prolactin. *J Clin Endocrinol Metab* 1988; 67: 460-4.
 20. Nestler JE. Sex hormone binding globulin. A marker for hyperinsulinemia and /or insulin resistance? *J Clin Endocrinol Metab* 1993; 76: 273-4.
 21. Adashi EY, Hseueh AJW, Yen SCC. Insulin enhancement of luteinizing hormone and follicle stimulating hormone release by cultured pituitary cells. *Endocrinology* 1981; 108: 1441-9.
 22. Nestler JE, Jakubowicz DJ. Decreases in ovarian cytochrome P450_{c17 α} activity and serum free testosterone after reduction of insulin secretion in polycystic ovary syndrome. *N Engl J Med* 1996; 335: 617-23.
 23. Dunaif A, Scott D, Finegood D, Quintana B, Whitcomb R. The insulin sensitizing agent troglitazone: a novel therapy for the polycystic ovary syndrome. *J Clin Endocrinol Metab* 1996; 81: 3299-306.
 24. Dunaif A, Graf M. Insulin administration alters gonadal steroid metabolism independent of changes in gonadotropin secretion in insulin-resistant women with polycystic ovary syndrome. *J Clin Invest* 1989; 83: 23-9.
 25. Dunaif A. Insulin resistance and ovarian dysfunction. In: Moller DE, editor. *Insulin resistance*. New York: John Wiley, 1993.
 26. Estelles A, Grancha S, Gilabert J. Abnormal expression of plasminogen activator inhibitor in patients with gestational trophoblastic disease. *Am J Pathol* 1996; 149: 1229-39.
 27. Halligan A, Bonnar J, Sheppard B. Haemostatic, fibrinolytic and endothelial variables in normal pregnancies and preeclampsia. *Br J Obstet Gynecol* 1994; 101: 488-92.
 28. Gris JC, Ripart-Neveu S, Maugard C. Respective evaluation of the prevalence of haemostasis abnormalities in unexplained primary early recurrent miscarriages. The Nimes Obstetricians and Haematologists (NOHA) Study. *Thromb Haemost* 1997; 77: 1096-103.
 29. Gris JC, Neveu S, Mares P. Plasma fibrinolytic activators and their inhibitors in women suffering from early recurrent abortion of unknown etiology. *J Lab Clin Med* 1993; 122: 606-15.
 30. Clifford K, Rai R, Watson H. Does suppressing luteinizing hormone secretion reduce the miscarriage rate? Results of a randomized controlled trial. *BMJ* 1996; 312: 1508-11.
-